

**VARIATIONS OF COMPOSITION OF ESSENTIAL OIL AND ACCUMULATION OF SALICYLIC ACID
IN *MENTHA ARVENSIS* VAR. *PIPERASCENS* (LAMIACEAE) WHEN INTRODUCED IN MOSCOW REGION**

© **O. V. Shelepova**,^{*} ¹**V. V. Kondrat'eva**,¹ **L. S. Olehnovich**,¹ **B. Ts. Zaitikh**,² **A. O. Ruzhitsky**²

¹ Tsitsin Main Botanical Garden RAS, Moscow, Russia

² Bach Institute of Biochemistry, Research Center of Biotechnology of the Russian Academy
of Sciences

*E-mail: shelepova-olga@mail.ru

REFERENCES

1. Clark G. S. 1998. Menthol. — *Perfumer et Flavorist*. 23(5): 33–46.
2. Burbott A. J., Loomis W. D. 1967. Effects of light and temperature on the monoterpenes of peppermint. — *Plant Physiol*. 42(1): 20–28.
3. Clark R. J., Menary R. C. 1980. Environmental effects on peppermint (*Mentha piperita* L.). I. Effect of day length, photon flux density, night temperature and day temperature on the yield and composition of peppermint oil. — *Austral. J. Plant Physiol*. 7(6): 685–692.
4. Santos V. M. C. S., Pinto M. A. S., Bizzo H., Deschamps C. 2012. Seasonal variation of vegetative growth, essential oil yield and composition of menthol mint genotypes at southern brazil. — *Biosci. J.* 28(5): 790–798.
5. Shelepova O. V., Kondratyeva V. V., Voronkova T. V., Olekhovich L. S. 2013. Correlation between the output and composition of essential oil and the level of salicylic acid in mint plants at different ontogenetic stages. — *Biology Bull.* 40(3): 275–280.
1. Fahlen A., Welander M., Wennersten R. 1999. Effects of light-temperature regimes on plant growth and essential oil yield of selected aromatic plants. — *J. Sci. Food Agric.* 73(1): 111–119.
2. Aflatuni A. 2005. The yield and essential oil content of mint (*Mentha* spp.) in Northern Ostrobothnia. Oulu. 52 p.
6. Tkachenko K. G. 2013. Essential oils families Apiaceae, Asteraceae and Lamiaceae in the Northwest Russia: Avtoref. dis. ... dokt. biol. nauk. St. Petersburg. 40 p. (In Russian)
7. Kondratyeva V. V., Kirichenko E. B., Safronova L. M., Voronkova T. V. 2000. Phytohormones of rhizomes of the mint of various geographic origin in its annual developmental cycle. — *Biology Bull.* 27(5): 471–476.
8. Yang J., Zhong J., Wang Z., Zhu Q., Liu L. 2002. Abscisic acid and cytokinins in the root exudates and leaves and their relationship to senescence and remobilization of carbon reserves in rice subjected to water stress during grain filling. — *Planta*. 215: 645–652.

9. Yuan S., Lin H.-H. 2008. Role of salicylic acid in plant abiotic stress. *Z. Naturforsch. C. Biosci.* 63(5–6): 313–320.
10. Scott M., Dat J. F. 1999. Salicylic acid and hydrogen peroxide in abiotic stress signalling in plants. — *Plant Physiol.* 39(3): 13–17.
11. Mahmoud S. S., Croteau R. B. 2003. Menthofuran regulates essential oil biosynthesis in peppermint by controlling a downstream monoterpene reductase. — *Proc. Nat. Acad. Sci. USA.* 100(24): 14481–14486.
12. Spinelli F., Cellini A., Marchetti L. 2011. Emission and function of volatile organic compounds in response to abiotic stress. In: *Abiotic stress in plants – mechanisms and adaptations*. Ed. by A. K. Shanker and B. Venkateswarlu. Rijeka. P. 367–394.
13. Singh A. K., Rain V. K., Naqvi A. A., Patra N. K., Birendra K., Ram P., Khanuja S. P. S. 2005. Essential oil composition and chemotarrays of menthol mint (*Mentha arvensis* L. f. *piperascens* Malinvaud ex Holmes) cultivars. — *Flavour Fragr. J.* 20(3): 302–305.